

TETANUS, DIPHTHERIA PERTUSSIS (Tdap) VACCINE

WHAT YOU NEED TO KNOW

1 Why get vaccinated?

Tdap (Tetanus, Diphtheria, Pertussis) vaccine can protect adolescents and adults against three serious diseases.

Tetanus, diphtheria, and pertussis are all caused by bacteria. Diphtheria and pertussis are spread from person to person. Tetanus enters the body through cuts, scratches, or wounds.

TETANUS (Lockjaw) causes painful tightening of the muscles, usually all over the body.

- It can lead to "locking" of the jaw so the victim cannot open his mouth or swallow. Tetanus leads to death in up to 2 cases out of 10.

DIPHTHERIA causes a thick covering in the back of the throat.

- It can lead to breathing problems, paralysis, heart failure, and even death.

PERTUSSIS (Whooping Cough) causes severe coughing spells, vomiting, and disturbed sleep.

- It can lead to weight loss, incontinence, rib fractures and passing out from violent coughing, pneumonia, and hospitalization due to complications.

In 2004 there were more than 25,000 cases of pertussis in the U.S. More than 8,000 of these cases were among adolescents and more than 7,000 were among adults. Up to 2 in 100 adolescents and 5 in 100 adults with pertussis are hospitalized or have complications.

2 Tdap and related vaccines

Vaccines for Adolescents and Adults

- **Tdap** was licensed in 2005. It is the first vaccine for adolescents and adults that protects against all three diseases.
- **Td** (tetanus and diphtheria) vaccine has been used for many years as booster doses for adolescents and adults. It does not contain pertussis vaccine.

Vaccines for Children Younger than 7 Years

- **DTaP** vaccine is given to children to protect them from these three diseases. Immunity can fade over time, and periodic "booster" doses are needed by adolescents and adults to keep immunity strong. (**DTP** is an older version of DTaP. It is no longer used in the United States.)
- **DT** contains diphtheria and tetanus vaccines. It is used for children younger than 7 who should not get pertussis vaccine.

3 Who should get Tdap vaccine and when?

Adolescents 11 through 18 years of age should get one booster dose of Tdap.

- A dose of Tdap is recommended for **adolescents who got DTaP or DTP as children** but have not yet gotten a dose of Td. The preferred age is 11-12.
- **Adolescents who have already gotten a booster dose of Td** are encouraged to get a dose of Tdap as well, for protection against pertussis. Waiting at least 5 years between Td and Tdap is encouraged, but not required.
- **Adolescents who did not get all their scheduled doses of DTaP or DTP** as children should complete the series using a combination of Td and Tdap.

Adults 19 through 64 years of age should substitute Tdap for one booster dose of Td. Td should be used for later booster doses.

- **Adults who expect to have close contact with an infant** younger than 12 months of age should get a dose of Tdap. Waiting at least 2 years since the last dose of Td is suggested, but not required.
- **Healthcare workers who have direct patient contact** in hospitals or clinics should get a dose of Tdap. A 2-year interval since the last Td is suggested, but not required.

An adolescent or adult who gets a severe cut or burn might need protection against tetanus infection. Tdap may be used if the person has not had a previous dose.

Td should be used rather than Tdap if Tdap is not available, and for:

- Anybody who has already gotten Tdap,
- Adults 65 years of age and older,
- Children 7 through 9 years of age.

If vaccination is needed during **pregnancy**, Td usually is preferred over Tdap. Ask your doctor. **New mothers** who have never received a dose of Tdap should get a dose as soon as possible after delivery.

Tdap may be given at the same time as other vaccines.

4 Some people should not get Tdap vaccine or should wait.

- Anyone who has had a **life-threatening allergic reaction** after a dose of DTP, DTaP, DT, or Td vaccine should not get Tdap.
- Anyone who has a **severe allergy to any component of the vaccine** should not get Tdap. Tell your health care provider if the person getting the vaccine has any known severe allergies.

continued ...

Talk with your doctor if the person getting the vaccine has a **severe allergy to latex**. Some Tdap vaccines should not be given to people with a severe latex allergy.

- Anyone who went into a **coma** or had a **long seizure** within 7 days after a dose of DTP or DTaP should not get Tdap, unless a cause other than the vaccine was found.
- Talk to your doctor if the person getting the vaccine:
 - has **epilepsy** or another **nervous system problem**,
 - had **severe swelling** or **severe pain** after a previous dose of any vaccine containing tetanus, diphtheria or pertussis,
 - has had **Guillain Barré Syndrome (GBS)**.

Anyone who has a **moderate or severe illness** on the day the shot is scheduled should usually wait until they recover before getting the vaccine. Those with a mild illness or low fever can usually be vaccinated.

5 What are the risks from Tdap vaccine?

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. However, the risk of a vaccine causing serious harm, or death, is extremely small.

If rare reactions occur with any new product, they may not be identified until many thousands, or even millions, of people have used the product. Like all vaccines, Tdap is being closely monitored for unusual or severe problems.

Clinical trials (testing before the vaccine was licensed) involved about 4,200 adolescents and about 1,800 adults. The following problems were reported. These are similar to problems reported after Td vaccine.

Mild Problems

(Noticeable, but did not interfere with activities)

- Pain (about 3 in 4 adolescents and 2 in 3 adults)
- Redness or swelling (about 1 in 5)
- Mild fever of at least 100.4°F (up to about 1 in 25 adolescents and 1 in 100 adults)
- Headache (about 4 in 10 adolescents and 3 in 10 adults)
- Tiredness (about 1 in 3 adolescents and 1 in 4 adults)
- Nausea, vomiting, diarrhea, stomach ache (up to 1 in 4 adolescents and 1 in 10 adults)
- Other mild problems reported include chills, body aches, sore joints, rash, and swollen lymph glands.

Moderate Problems

(Interfered with activities, but did not require medical attention)

- Pain at the injection site (about 1 in 20 adolescents and 1 in 100 adults)
- Redness or swelling (up to about 1 in 16 adolescents and 1 in 25 adults)
- Fever over 102°F (about 1 in 100 adolescents and 1 in 250 adults)
- Nausea, vomiting, diarrhea, stomach ache (up to 3 in 100 adolescents and 1 in 100 adults)
- Headache (1 in 300)

Severe Problems

(Unable to perform usual activities; required medical attention)

- None were seen among adolescents.
- In the adult clinical trial, two adults had nervous system problems after getting the vaccine. These may or may not have been caused by the vaccine. They went away on their own and did not cause any permanent harm.
- A severe allergic reaction could occur after any vaccine. They are estimated to occur less than once in a million doses.

A person who gets these diseases is much more likely to have severe complications than a person who gets Tdap vaccine.

6 What if there is a severe reaction?

What should I look for?

- Any unusual condition, such as a high fever or behavior changes. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?

- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form.

Or you can file this report through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS does not provide medical advice.

7 The National Vaccine Injury Compensation Program

In the event that you or your child has a serious reaction to a vaccine, a federal program has been created to help pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call 1-800-338-2382 or visit their website at www.hrsa.gov/vaccinecompensation.

8 How can I learn more?

- Ask your immunization provider. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO)
 - Visit CDC's National Immunization Program website at www.cdc.gov/nip



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL CENTER FOR IMMUNIZATION AND RESPIRATORY DISEASES

TETANUS AND DIPHTHERIA VACCINE (Td)

What you need to know before you or your child gets the vaccine



U.S. DEPARTMENT OF HEALTH & HUMAN SERVICES
Public Health Service
Centers for Disease Control
and Prevention

ABOUT THE DISEASES

Tetanus (lockjaw) and diphtheria are serious diseases. Tetanus is caused by a germ that enters the body through

a cut or wound. Diphtheria spreads when germs pass from an infected person to the nose or throat of others.

Tetanus causes:
serious, painful spasms of all muscles

It can lead to:
- "locking" of the jaw so the patient cannot open his or her mouth or swallow

Diphtheria causes:
a thick coating in the nose, throat, or airway

It can lead to:
- breathing problems
- heart failure
- paralysis
- death

ABOUT THE VACCINES

Benefits of the vaccines

Vaccination is the best way to protect against tetanus and diphtheria. Because of vaccination, there are many fewer cases of these diseases. Cases are rare in children because most get DTP (Diphtheria, Tetanus, and Pertussis), DTaP (Diphtheria, Tetanus, and acellular Pertussis), or DT (Diphtheria and Tetanus) vaccines. There would be many more cases if we stopped vaccinating people.

When should you get Td vaccine?

Td is made for people 7 years of age and older.

People who have not gotten at least 3 doses of any tetanus and diphtheria vaccine (DTP, DTaP, or DT) during their lifetime should do so using Td. After a person gets the third dose, a Td dose is needed every 10 years all through life.

Other vaccines may be given at the same time as Td.

Tell your doctor or nurse if the person getting the vaccine:

- ever had a serious allergic reaction or other problem with Td, or any other tetanus and diphtheria vaccine (DTP, DTaP, or DT)
- now has a moderate or severe illness
- is pregnant

If you are not sure, ask your doctor or nurse.

What are the risks from Td vaccine?

As with any medicine, there are very small risks that serious problems, even death, could occur after getting a vaccine.

The risks from the vaccine are much smaller than the risks from the diseases if people stopped using vaccine.

Almost all people who get Td have no problems from it.

Mild problems

If these problems occur, they usually start within hours to a day or two after vaccination. They may last 1-2 days:

soreness, redness, or swelling where the shot was given

These problems can be worse in adults who get a vaccine very often.

Acetaminophen or ibuprofen (non-aspirin) may be used to reduce soreness.

Severe problems

These problems happen very rarely:

serious allergic reaction

deep, aching pain and muscle wasting in upper arm(s). This starts 2 days to 4 weeks after the shot, and may last many months.

What to do if there is a serious reaction:

- ☒ Call a doctor or get the person to a doctor right away.
- ☒ Write down what happened and the date and time it happened.
- ☒ Ask your doctor, nurse, or health department to file a Vaccine Adverse Event Report form or call: (800) 822-7967 (toll-free)

The National Vaccine Injury Compensation Program gives compensation (payment) for persons thought to be injured by vaccines. For details call:

(800) 338-2382 (toll-free)

If you want to learn more, ask your doctor or nurse. She/he can give you the vaccine package insert or suggest other sources of information.

